

## On the volume and the Chern-Simons invariant for the 2-bridge link orbifolds

*Ji-Young Ham*

*Department of Science, Hongik University, Seoul, Korea,  
Department of Mathematical Sciences, Seoul National University, Seoul, Korea  
jyoungham1@gmail.com*

This is joint work with Joongul Lee, Alexander Mednykh, and Aleksey Rasskazov

We extend some part of the unpublished paper [1] written by Mednykh and Rasskazov. Using the approach indicated in this paper we derive the Riley–Mednykh polynomial for some family of the two bridge link orbifolds. As a result we obtain explicit formulae for the volume of cone–manifolds and the Chern–Simons invariant of orbifolds of the knot with Conway’s notation  $C(2n, 4)$ .

### References

- [1] A. Mednykh, A. Rasskazov, On the structure of the canonical fundamental set for the 2–bridge link orbifolds. (1998).
- [2] A. Mednykh, A. Rasskazov, Volumes and degeneration of cone-structures on the figure–eight knot. *Tokyo J. Math.* **29(2)** (2006) 445–464.
- [3] Ji-Y. Ham, A. Mednykh, V. Petrov, Trigonometric identities and volumes of the hyperbolic twist knot cone-manifolds. *J. Knot Theory Ramifications* **23(12)** (2014)1450064, 16.
- [4] Ji-Y. Ham, J. Lee, Explicit formulae for Chern–Simons invariants of the twist knot orbifolds and edge polynomials of twist knots. *arXiv:1411.2383* (2014). Preprint.
- [5] Ji-Y. Ham, J. Lee, The volume of hyperbolic cone–manifolds of the knot with Conway’s notation  $C(2n, 3)$ . *arXiv:1512.05481* (2015). Preprint.
- [6] Ji-Y. Ham, J. Lee, Explicit formulae for Chern–Simons invariants of the hyperbolic orbifolds of the knot with Conway’s notation  $C(2n, 3)$ . *arXiv:1601.00723* (2016). Preprint.